



GENERAL APPLICATION GUIDE

The following are only suggested treatment protocols developed and provided by the licensed practitioner community. Each patient should be consulted and appropriate treatment offered for the specific indications by a licensed practitioner. Please administer in accordance with current FDA regulations.

Cosmetic/Dermatology

| | | | |
|------------------|------------|------|--|
| Acne | RICHGEN I | 1 ml | topically 100% |
| Burns | RICHGEN II | 2 ml | topical spray 100% |
| CO2 lasering | RICHGEN I | 1 ml | topically 100% post laser |
| Hair restoration | RICHGEN II | 2 ml | with 5 ml saline per 1 ml product (injection with 3/16 th needle) |
| Micro-needling | RICHGEN I | 1 ml | topically 100% post micro-needling |
| Scar keloid | RICHGEN I | 1 ml | with 1 ml saline injected locally and topically |
| Surgery | RICHGEN I | 1 ml | topically 100%, membrane can be inserted before closure |

Dental

| | | | |
|--------------------------|-----------|------|--|
| Implants, bone, Canal | RICHGEN I | 1 ml | with 1 ml saline injected to area and around |
|--------------------------|-----------|------|--|

Endocrinology

| | | | |
|------------------|------------|------|-------------|
| Immune Disorders | RICHGEN II | 2 ml | with saline |
| Lyme | RICHGEN II | 2 ml | with saline |

Neuropathy

Diabetes RICHGEN II 3 ml-2 ml with saline, 1 ml with 1 ml saline injection and topically to wound

OBGYN

Pelvic floor RICHGEN I 1 ml 100% with 1 ml saline injected to area
Pudental nerve RICHGEN I 1 ml 100% with 1 ml saline injected to area
P or O (G spot) RICHGEN I 1 ml 100% with 1 ml saline injected area around & to area
RF FemiLift RICHGEN I 1 ml 100% with 1 ml saline injected to O (G) Spot 3 days later

Oncology

Cancer, tumors RICHGEN II There are not enough studies yet

Ophthalmology

Macular degeneration RICHGEN combo 1 ml with 2 ml saline in an eye dropper to administer one drop every 15 seconds for 2 hours for one eye.
If both eyes, double the amount.
3 ml with ample saline by push.
Product needs to be used quickly. We have .5 ml vials in stock sometimes.
If not, order two weeks in advance so that the product can be thawed in smaller amounts to keep it fresh during the 2-hour process.

Orthopedic

Bone, Tendon RICHGEN I 1 ml 100% or with 1 ml saline injected per area
Spine: RICHGEN I 1 ml 100% or with 1 ml saline injected per vertebra, disc area

2 ml suggested for large body mass/tall patients. Plus, local vein injection used to rebuild vein connections for paralyzed patients. Injection for disc regeneration is same as injecting cortisone.
See Ortho [Application Guide](#) for knees and joints, pain, bone regeneration.

Podiatry

See Ortho [Application Guide](#) for foot and ankle.

Pulmonary/Respiratory

| | | | |
|--------------|------------|---------|--|
| Asthma, COPD | RICHGEN II | 1-2 ml | 50% with 1 ml saline each IM tricep injection each arm |
| | | 2 – 3ml | Optional: nasal passage injection with 1:1 saline after numbing (Formerly +25% by mist/neb with 1 - 1.5 ml saline was used and is being replaced with IM tricep.) |

Rheumatology

| | | | |
|--------------------|------------|-----------|----------------------------------|
| Inflammation, pain | RICHGEN II | 2 ml | 1 – 1.5 ml with saline |
| | | .5 – 1 ml | injected to finger or toe joints |

Urologist

| | | | |
|---------------|------------|------|------------------------------------|
| Kidneys | RICHGEN II | 2 ml | 100% with 4-5 ml saline per kidney |
| P (ED) | RICHGEN I | 1 ml | 100% injection with 1 ml saline |
| O (G spot) | RICHGEN I | 1 ml | 100% injection with 1 ml saline |
| Prostatectomy | RICHGEN II | 2 ml | 100% with 4-5 ml saline injected |

Each patient responds differently based on a variety of factors. The RichSource product line represents – RICHGEN – that remains consistent in quality and preservation protocols for each application. Being undifferentiated pluripotent results are beyond limited multipotent cord blood that is not suitable for most applications contrary to marketing. Not all disorders are listed. The above suggestions are not to be interpreted or relied upon as a medical diagnosis, treatment protocol as each patient may require custom applications. Multiple treatments may be required. Each treatment for a patient is under a physician consultation and recommendation independent of this website content and products or representatives and should be in accordance with FDA current

regulations.

The following are not allowed for three days prior and for 8 weeks after injection treatments:

No Aspirin No Advil No Aleve

No Ibuprofen No Naproxen No Icing

Have questions? RichSource has area executives who will personally [call on you](#).

SAFER UNRIVALED UNPRECEDENTED WORLDWIDE