



HIP AND SACROILIAC SIJ APPLICATION GUIDE

The following are only suggested treatment protocols developed and provided by the licensed practitioner community. Each patient should be consulted and appropriate treatment offered for the specific indications by a licensed practitioner. Please administer in accordance with current FDA regulations or Health Ministry regulations of your country. *

HIP

RICHGEN I or II

Use 1 ml or 2ml depending on the condition of the patient with 1:2 or 1:3 saline for size of patient.

First, shockwave the area to breakup mineralization and exercise the joint to increase inflammation.

Use sterile gel on the skin with ultrasound, C Arm or other visual aid to guide the injections.

Next, for anterior injections look for the dark cavity area to inject.

For posterior injections lay the patient on their side and flex the knee. Look for the dark, small crescent shaped cavity to inject.

Guide the needle into the dark spot(s) where needed.

Lateral injection is not advised.

SACROILIAC SIJ

RICHGEN I or II

Use 1 ml or 2ml depending on the condition and size of the patient with 1:8 or 1:10 saline as this joint area is large to fill.

First, shockwave the area to breakup mineralization and exercise the joint to increase inflammation.

Use sterile gel on the skin with ultrasound, C Arm or other visual aid to guide the injections.

*Each patient responds differently based on a variety of factors. The RichSource product line represents – RICHGEN – that remains consistent in quality and preservation protocols for each application. Being undifferentiated pluripotent results are beyond limited multipotent cord blood that is not suitable for most applications contrary to marketing. Not all disorders are listed. The above suggestions are not to be interpreted or relied upon as a medical diagnosis, treatment protocol as each patient may require custom applications. Multiple treatments may be required. Each treatment for a patient is under a physician consultation and recommendation independent of this website content and products or representatives.

The following are not allowed for three days prior and for 8 weeks after injection treatments:

No Aspirin No Advil No Aleve

No Ibuprofen No Naproxen No Icing

RICHGEN contains 3% DMSO (dimethyl sulfoxide).