

## Hair Restoration Application Guide



Before RICHGEN Treatment



5 Weeks After Using RICHGEN

**RICHGEN I** or **II** depending on the size of the area to be treated.  
One ml for partial area or 2 ml for the entire scalp.

Use 10 ml of saline per 1 ml of RICHGEN.

Use only 3/16th length needle 30 gauge with 1 ml syringe used for insulin.

Thaw the RICHGEN vial(s) in the palm of your gloved hand for three (3) minutes.

Carefully place RICHGEN 1 ml with the 10 ml saline in one 12 ml + syringe.

Then extract fill one dozen 1 ml syringes with the combined fluid. Extract all the RICHGEN cells from the vials with some of the saline after emptied to remove all the cells from the vial.

Repeat the process with second ml if using 2 ml.

After the triple lidocaine numbing crème has taken effect in 30 minutes, proceed with the injections approximately 5 – 7mm apart. If additional pain medication is needed, inject lidocaine into the scalp.

It is best if the patient does not wash their scalp for 24 – 48 hours. When shampooing, it is best to use very, very gentle fingertip light massaging of the shampoo through the hair and rinse, for a month. An additional treatment may be needed in two months.

**Supplies Provider:**

Harry Schein, Inc. 800.772.4346, press # for sales 8am-8:30pm EST or

Ask for Earl at 626.688.4518

Steriject #1734233 30 gauge 3/16 needles 100/box

Becton Dickinson Syringes #9875471 1 ml 100/box

Each patient responds differently based on a variety of factors. The RichSource product line represents – RICHGEN – that remains consistent in quality and preservation protocols for each application. Being undifferentiated pluripotent results are beyond limited multipotent cord blood that is not suitable for most applications contrary to marketing. RICHGEN is richer than PRP. The above suggestive application guide is not to be interpreted or relied upon as a medical diagnosis or treatment protocol as each patient may require custom applications. Multiple treatments may be required. Each treatment for a patient is under a physician consultation and recommendation independent of this website content and products or representatives. The following are not allowed for three days prior and for 8 weeks after injection treatments:

No Aspirin No Advil No Aleve

No Ibuprofen No Naproxen No Icing

Have questions? RichSource has area executives who will personally [call on you](#).

RICHGEN contains 3% - 5% DMSO.

**WARNING: Patients taking heart medications and patients who are allergic to sulfur may be at risk if treated.**