



RICHSOURCE®

STEM CELLS, INC

The following are only suggested treatment protocols developed and provided by the licensed practitioner community and provided for information purposes only between peers. Each patient should be consulted and appropriate treatment offered for the specific indications by a licensed practitioner. Please administer in accordance with current FDA guidelines and regulations.

Tendon-Fascia Application Guide

Stimulate inflammatory signaling.

Radial pulse therapy, ESWT, needling, walking, surgery.

Think of anatomy, consider vascular supply to the tendon or fascia.

Inject along the paratenon, the covering of the tendon, or around the fascia, so RICHGEN allograft is taken up into microcirculation of the tendon.

Use imaging but you do not need to go into the tendon or fascia mass.

Inject peri tendon versus into tendon or fascia.

Active rehabilitation, protect if necessary but do not immobilize.

Use RICHGEN I or II - 1 or 2 ml depending on area and acuteness with 1:1 saline. Then allow 4 weeks for the regeneration to work. Repeat if needed.

Each patient responds differently based on a variety of factors. The RichSource product line represents – RICHGEN – that remains consistent in quality and preservation protocols for each application. Being undifferentiated pluripotent results are beyond limited multipotent cord blood that is not suitable for most applications contrary to marketing. Not all disorders are listed. The above suggestions are not to be interpreted or relied upon as a medical diagnosis, treatment protocol as each patient may require custom applications. Multiple treatments may be required. Each treatment for a patient is under a physician consultation and recommendation independent of this website content and products or representatives and should be in accordance with FDA current guidelines and regulations.

The following are not allowed for three days prior and for 8 weeks after injection treatments:

No Aspirin No Advil No Aleve

No Ibuprofen No Naproxen No Icing

RICHGEN contains 3% DMSO (dimethyl sulfoxide).